

Monochlorobenzene Contaminated Site Characterization by the Use of ^{37}Cl , ^{13}C and ^2H -Compound-Specific Isotope Analysis (CSIA), Biological Molecular Techniques (BMTs) and Numerical Modeling

Massimo Marchesi (massimo.marchesi@polimi.it), Ilaria Pietrini, Matteo Antelmi and Luca Alberti (luca.alberti@polimi.it) (Politecnico di Milano, Milano, Italy)
Tatiana Stella and Andrea Franzetti (Università Statale di Milano-Bicocca, Milano, Italy)
Danilo Antonelli and Francesca de Ferra (ENI Group, Milano, Italy)
Ramon Aravena, (University of Waterloo, Waterloo, ON, Canada)
Orfan Shouakar-Stash (orfan@it2isotopes.com) (Isotope Tracer Technology Inc., Canada)

Background/Objectives. Chlorinated solvent contaminated sites entail costly management costs and often create significant impacts to the environment. Innovative tools such as compound-specific isotope analysis (CSIA), molecular biological techniques (MBTs) and fate and transport models can provide unequivocal information for site characterization and assessment of natural attenuation processes, particularly biodegradation. This contribution explores the potential for the tri-element CSIA approach, MBTs and transport models to study a heavy monochlorobenzene (MCB) and benzene contamination at a mega-site where potential multiple sources are present.

Approach/Activities. Microcosm experiments were developed to estimate enrichment factors for ^{13}C and ^{37}Cl , and to investigate specific biomarkers under different conditions (soil slurry, natural and stimulated, a- and anaerobic). Field investigations were performed including $d^{37}\text{Cl}$, $d^{13}\text{C}$ and $d^2\text{H}$ for MCB and benzene and illumina sequencing and qPCR coupled to reactive transport modeling efforts.

Results/Lessons Learned. A significant difference in the $d^{13}\text{C}/d^2\text{H}$, $d^{13}\text{C}/d^{37}\text{Cl}$, $d^{37}\text{Cl}/d^2\text{H}$ and $d^{13}\text{C}/d^{37}\text{Cl}/d^2\text{H}$ for MCB and $d^{13}\text{C}/d^2\text{H}$ for benzene - in the high concentration spots - allowed distinguishing two distinct sources and plumes (merged at some areas). By the use of MNWs a detailed vertical distribution of the contaminants through the aquifer was assessed. Concerning natural attenuation processes, although the site is characterized by reducing conditions with high concentrations of CH_4 , MBTs results indicated the presence of mainly aerobic potential degraders within the indigenous community at the site and ruled out a significant presence of dehalococoides or dehalobacters. Microcosms experiments and soil slurry with in-situ cultures showed a unique $d^{13}\text{C}/d^{37}\text{Cl}$ trend with almost no enrichment for ^{13}C but a ^{37}Cl ? of - 0.7 ‰.

The dual $d^{13}\text{C}/d^{37}\text{Cl}$ data also seem to indicate aerobic biodegradation processes as the main natural attenuation mechanism. Finally, regarding benzene, the $d^{13}\text{C}/d^2\text{H}$ data showed that benzene present at the site is probably not linked to MCB dehalogenation processes but is rather a primary contaminant; these data also showed benzene is being attenuated by biodegradation. Since not varying as a consequence of biodegradation processes, $d^{13}\text{C}$ and $d^2\text{H}$ for MCB were used in combination with the transport models to estimate the contribution to the contamination from the two distinct sources. After estimating the mixing ratios among the plumes, $d^{37}\text{Cl}$ was successfully used as a tracer to assess the natural attenuation processes at the site.