

Treatment of Benzene Contamination Using Rhizoremediation at a Petrochemical Facility in Brazil

Flavio Coelho (flavio.coelho@erm.com) (ERM Brasil, Rio de Janeiro, RJ, Brazil)

Background/Objectives. Phytoremediation, the use of plants to degrade toxic contaminants in the environment, involves a number of processes including phytoextraction, phytotransformation, phytostabilization, phytovolatilization and rhizofiltration. While some success has been reported using plants alone in bioremediation, the use of plants in conjunction with plant associated bacteria offers a lot of potential. Rhizoremediation systems rely on a synergistic relationship between suitable plants and their root associated microbial communities, when degradation is facilitated through a rhizosphere effect where plants exude organic compounds through their roots and thereby increase the density and activity of potential hydrocarbon degrading microorganisms in the area surrounding the roots. The chemical facility is located at the banks of the Guanabara Bay, in Rio de Janeiro, at an environmentally protected area. The existing benzene contamination impacts an area of approximately 120,000 sq. meters, in concentrations of up to 300 mg/L. This approach will be used in areas where concentrations are below 10 mg/L, which comprises approximately 90% of the plume area.

Approach/Activities. A bench-scale pilot test was performed to determine the potential for in situ bioremediation at the site. The test was designed to evaluate the effectiveness of anaerobic and aerobic ISB to treat the benzene. The test concluded that benzene-degrading bacteria existed at the soil and groundwater at the site, and that oxygenation of the media favored the reduction of the VOCs. Another study is being conducted to determine, between the several native plants to the region, which ones will be more efficient on the overall objective of the project, which is to slow down the groundwater flow towards the bay and at the same time provide the ideal rhizosphere to enhance bacterial and fungal activity and degrade the VOCs.

Results/Lessons Learned. The treatability tests indicated a removal efficiency 99.8% or higher reductions in benzene and complete reductions of toluene, ethylbenzene, and total xylenes. Oxygenation of the sand soil and groundwater would provide for greater and more rapid reductions in VOCs. The pilot test activities will be conducted in December 2015 to March 2016 and this paper will present the treatability tests and the pilot test results with special attention to the performance and implementation of the field programs, challenges imposed by the Site and technology conditions and optimizations applied to address the overall performance of the strategy. Full-scale field work implementation is scheduled to start in November 2016. The paper will also include the environmental agency response to the proposed rhizoremediation technology and the strategy used to convey the technology both to the agency and the communities around the site.